

Research Paper

**Cite this article:** Menasria A, Barčák D, Kaouachi N, Bensouilah M, Scholz T, Hernández-Orts JS (2019). Redescription of *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) (Acanthocephala: Quadrigyridae), a parasite of the Algerian barb *Luciobarbus callensis* (Valenciennes) (Cyprinidae) in Algeria, and first molecular data. *Journal of Helminthology* 1–12. <https://doi.org/10.1017/S0022149X19000737>

Received: 13 May 2019

Revised: 16 July 2019

Accepted: 19 July 2019







**Key words:**

Acanthocephala; Eoacanthocephala; freshwater fish; taxonomy; 28S rDNA; phylogeny; North Africa

**Author for correspondence:**

J.S. Hernández-Orts, E-mail: [jesus.hernandez.orts@gmail.com](mailto:jesus.hernandez.orts@gmail.com)

# Redescription of *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) (Acanthocephala: Quadrigyridae), a parasite of the Algerian barb *Luciobarbus callensis* (Valenciennes) (Cyprinidae) in Algeria, and first molecular data

A. Menasria<sup>1</sup> , D. Barčák<sup>2</sup> , N. Kaouachi<sup>1</sup> , M. Bensouilah<sup>3</sup> , T. Scholz<sup>4</sup>   
and J.S. Hernández-Orts<sup>5</sup> 

<sup>1</sup>Laboratory of Terrestrial and Aquatic Ecosystems, Faculty of Natural and Life Sciences, University of Mohamed Cherif Messaadia, BP 1553, Annaba Road, 41000 Souk Ahras, Algeria; <sup>2</sup>Institute of Parasitology, Slovak Academy of Sciences, Hlinkova 3, 040 01 Košice, Slovak Republic; <sup>3</sup>Ecobiology Laboratory for Marine Environments and Coastal Areas, Faculty of Sciences, University of Badji Mokhtar, BP 12 El-Hadjer, 23000 Annaba, Algeria; <sup>4</sup>Institute of Parasitology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, 370 05 České Budějovice, Czech Republic and <sup>5</sup>Centro de Investigación Aplicada y Transferencia Tecnológica en Recursos Marinos Almirante Storni (CIMAS – CCT CONICET – CENPAT), Güemes 1030, 8520 San Antonio Oeste, Río Negro, Argentina

## Abstract

*Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951), an insufficiently described quadrigyrid acanthocephalan of cyprinid fishes from Northwest Africa, is redescribed based on recently collected specimens from the Algerian barb *Luciobarbus callensis* (Valenciennes) in Algeria. Newly observed morphological features for *A. (A.) maroccanus* include the arrangement of proboscis hooks (not in regular circles), the male reproductive structures extending into the copulatory bursa and the presence of a para-receptacle structure and vaginal sleeve. The mechanism of copulation of this acanthocephalan is described based on several copulating pairs. The phylogenetic position of *A. (A.) maroccanus* within Eoacanthocephala was assessed based on partial 28S rDNA sequences. Maximum likelihood and Bayesian inference analyses placed *A. (A.) maroccanus* in a clade with *Palliolisentis (Demidueterospinus) ophiocephalus* (Thapar, 1931), both species included in the Quadrigyridae, the only family within the Gyraacanthocephala.

## Introduction

*Acanthogyrus* Thapar, 1927 is a cosmopolitan genus of quadrigyrid acanthocephalans found in freshwater and marine fish (Amin, 2005). *Acanthogyrus* is subdivided in two subgenera: *Acanthogyrus* Thapar, 1927 with two species and *Acanthosentis* Verma & Datta, 1929 with 45 known species (Amin, 2005, 2013; Amin *et al.*, 2017). These subgenera are distinguished by the number of hooks on the proboscis: 24 in *Acanthogyrus* (three circles of eight hooks each) and 18 in *Acanthosentis* (three circles of six hooks each) (Amin & Hendrix, 1999).

*Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) was described from *Luciobarbus setivimensis* (Valenciennes) (syn. *Barbus setivimensis* Valenciennes) from Azrou, Morocco by Dollfus (1951). The species was described only briefly and illustrated poorly by Dollfus (1951) and Meddour *et al.* (2010); the latter authors reported the species from the Algerian barb *Luciobarbus callensis* (Valenciennes) (syn. *Barbus callensis* Valenciennes) in Algeria. Recently, we examined newly collected material of *A. (A.) maroccanus* from *L. callensis* from Algeria. In the present paper, *A. (A.) maroccanus* is redescribed and scanning electron microscopical (SEM) micrographs are also provided. Our observations reveal some morphological features not reported in previous morphological descriptions of this taxon. In addition, we provide an estimation of the phylogenetic framework, based on partial sequences for the large subunit of the ribosomal RNA gene (28S rDNA), to test the relationships of the genus *Acanthogyrus* (order Gyraacanthocephala) within the class Eoacanthocephala.

## Materials and methods

### Sample collection

A total of 359 *L. callensis* were collected from three localities in Algeria between March 2016 and March 2018: (i) Oued Charef dam lake, Sedrata, Souk Ahras City (36°42'00"N, 7°23'00"E) ( $n = 289$ ; 26.5–49.0 cm in total length); (ii) Zit Emba dam lake, Skikda City (36°44'55"N,

7°23'08"E) ( $n = 36$ ; 19.0–39.0 cm in total length); and (iii) Beni Haroun dam lake, Mila City (36°33'49"N, 6°16'14"E) ( $n = 34$ ; 30.0–40.0 cm in total length). The fish were captured with a net, transported alive to the laboratory and euthanized by spinal severance. Acanthocephalans found were placed in Petri dishes with tap water, left in a refrigerator for 1–15 h and then fixed in 4% formaldehyde solution (=formalin) or in 96% ethanol.

### Morphological description

Selected acanthocephalans fixed in 4% formalin were gently punctured with a fine needle and stained with Mayer's pararcarmine, washed in distilled water, dehydrated in ethanol, cleared in methyl salicylate and mounted as permanent slides in Canada balsam. Mounted acanthocephalans were examined using an Olympus BX51 microscope (Olympus Corporation, Tokyo, Japan). Measurements, taken using the QuickPhoto Micro Microscope Software (Promicra, Prague, Czech Republic), are given in micrometres ( $\mu\text{m}$ ) unless otherwise stated, and are expressed as the range, with the mean and the number of measurements in parentheses. Fully mature eggs were measured from pictures of eggs *in situ* through the body wall of female worms. Infection parameters were estimated following Bush *et al.* (1997).

Eight adult acanthocephalans (four males and four females), fixed in 4% formalin, were studied with SEM. Worms were post-fixed in 2% osmium tetroxide for 2 h, washed in 0.1 M phosphate buffer, dehydrated through an acetone series, critical point-dried and sputter coated with gold. Samples were examined using a JEOL JSM 7401-F scanning electron microscope at an accelerating voltage of 4 kV in the Laboratory of Electron Microscopy, Institute of Parasitology, Biology Centre ASCR, České Budějovice, Czech Republic.

Voucher specimens are deposited in the Helminthological Collection of the Institute of Parasitology (IPCAS), Biology Centre ASCR, České Budějovice, Czech Republic; the Helminthological Collection of the Museo de La Plata (HCMLP-He), Buenos Aires, Argentina; and the Natural History Museum (NHMUK), London, UK.

### Molecular analysis

Genomic DNA was isolated from three ethanol-fixed specimens by DNeasy® Blood & Tissue Kit (QIAGEN, Hilden, Germany) following manufacturer's instructions, except for the last step, in which elution buffer was replaced by deionized water. The target D1–D3 region of 28S rDNA was amplified by using LSU5 and 1500R primers (Littlewood *et al.*, 2000; Olson *et al.*, 2003). The amplification was performed as (i) denaturation at 95°C for 3 min; (ii) 40 cycles of 94°C for 30 s, 55°C for 40 s, 72°C for 90 s; and (iii) termination at 72°C for 5 min. The polymerase chain reaction (PCR) products were verified on 1% agarose gel and enzymatically purified by exonuclease I and shrimp phosphatase (Werle *et al.*, 1994). Sanger sequencing was held at GATC Biotech (Cologne, Germany) using both amplification primers and 900F internal primer (Olson *et al.*, 2003). Contiguous sequence was submitted to the GenBank database.

Newly generated sequences were aligned together with sequence data for species included in the class Eoacanthocephala in order to assess the phylogenetic relationships of the genus *Acanthogyryus* within this class. The dataset for the Eoacanthocephala comprised sequences for four genera and 18 species included in the order Neoecchinorhynchida, and a single sequence of *Palliolisentis*

(*Demidueterospinus*) *ophiocephalus* (Thapar, 1931) of the order Gyraacanthocephala, as currently available in the GenBank database (see table 1 for details). The partial sequences of the 28S rRNA gene of *Acanthogyryus* (*Acanthosentis*) *tilapiae* Baylis, 1947 (*nec* 1948; see Baylis, 1947) available in the GenBank database (ATU53000) generated by Chenuil *et al.* (1997) were not included in the analysed dataset because of the very short length of the sequence (311 bp). Based on previous molecular phylogenies, sequences of species included in the classes Archiacanthocephala, Palaeacanthocephala and Polyacanthocephala were used as external groups (table 1). Contiguous sequences were assembled and inspected for errors in platform Geneious® v10.1.3 (Kearse *et al.*, 2012), and gene alignments were built by the program ClustalW (Thompson *et al.*, 1994), which is implemented in the website <http://www.genome.jp/tools/clustalw/>, with the approach 'SLOW/ACCURATE' and weight matrix 'CLUSTALW (FOR DNA)'. Maximum likelihood analysis was performed by RAXML v8.2.11 (Stamatakis, 2014) with 1000 bootstrap replicates. A Bayesian inference tree was generated by MrBayes v3.2.2 (Huelsenbeck & Ronquist, 2001), in which four independent MC3 runs of ten million generations each were realized, and for each run, two chains were used; tree topologies were sampled every 1000 generations, the heating parameter value was 0.2 and 'Burn-in' was set to 25%. The evolution model GTR+G for both RAXML and MrBayes was estimated with the program jModelTest v2.1.10 (Darriba *et al.*, 2012) using corrected Akaike Information Criterion. Genetic distances (uncorrected p-distance) and the number of parsimony informative sites in the alignment were calculated with Geneious® and PAUP 4.0a165, respectively (Swofford, 2002).

### Results

#### *Acanthogyryus* (*Acanthosentis*) *maroccanus* (Dollfus, 1951) (figs 1–4; supplementary table S1)

#### Redescription

*General.* Quadrigyridae, with characters of *Acanthogyryus* and subgenus *Acanthosentis*. Sexual dimorphism evident, females larger than males (fig. 1a, c). Proboscis short, cylindrical, with small apical organ, pump in middle, armed with three irregularly arranged circles (especially anterior and middle circle) of six rooted hooks each, and one mononucleated large cell (figs 1b, d, 2a, 3a, b and 4a, b). Two lateral proboscis hooks in anterior and middle circle displaced posteriorly, but do not differ in size from two dorsal and two ventral hooks of each circle (figs 2a, 3b, c and 4b). Anterior hooks of similar length as middle hooks (fig. 2a). Posterior hooks shorter than anterior and middle hooks (figs 3b, d and 4b). Roots simple, without manubria, shorter than blades (fig. 2a). Neck present, unarmed (figs 3b and 4b). Trunk short, cylindrical (fig. 1a, c). Giant hypodermal nuclei 9–12; 2–3 ventral, 6–9 dorsal (fig. 1a), not observed in most specimens (especially in females). Complete circles of spines on anterior trunk scarce, usually fewer spines dorsally (figs 1b, d, 3a and 4a). Spines become widely spaced with no or fewer dorsal spines posteriorly (figs 1b, d, 3a and 4a). Proboscis receptacle single-walled (fig. 1b, d), 2.0–5.5 (3.9;  $n = 4$ ) times longer than proboscis in males and 2.9–5.7 (4.1;  $n = 3$ ) times longer in females, with small, ellipsoidal cerebral ganglion close to its base (fig. 1b). Para-receptacle structure evident in both sexes, contiguous with ventral side of receptacle (fig. 1b, d). Lemnisci

**Table 1.** Acanthocephalans included in the phylogenetic analyses with data on the host, locality and GenBank accession number (28S rDNA).

Species	Host	Locality	GenBank accession no.	Reference
<b>Class: Eoacanthocephala</b>				
Order: Gyracanthocephala				
Family: Quadrigyridae				
Genus: <i>Acanthogyrus</i>				
<i>A. (A.) maroccanus</i>	<i>L. callensis</i>	Souk Ahras City, Algeria	MK953673	Present study
Genus: <i>Palliolisentis</i> Machado Filho, 1960				
Subgenus: <i>Demidueterospinus</i> Amin, Heckmann, Ha, Luc & Doanh, 2000				
<i>P. (D.) ophiocephalus</i> (Thapar, 1931)	<i>Channa punctata</i> (Bloch)	India	KF700099	Unpublished
Order: Neoechinorhynchida				
Family: Neoechinorhynchidae				
Genus: <i>Atactorhynchus</i> Chandler, 1935				
<i>A. duranguensis</i> Salgado-Maldonado, Aguilar-Aguilar & Cabañas-Carranza, 2005	<i>Cyprinodon meeki</i> Miller	Durango, Mexico	KY077080	Pinacho-Pinacho <i>et al.</i> (2017)
Genus: <i>Floridosentis</i> Ward, 1953				
<i>F. mugilis</i> (Machado Filho, 1951)	<i>Mugil cephalus</i> L.	Veracruz, Mexico	JQ436495	Rosas-Valdez <i>et al.</i> (2012)
<i>F. pacifica</i> Bravo-Hollis, 1969	<i>Mugil curema Valenciennes</i>	Guerrero, Mexico	JQ436531	Rosas-Valdez <i>et al.</i> (2012)
Genus: <i>Mayarhynchus</i> Pinacho-Pinacho, Hernández-Orts, Sereno-Urbe, Pérez-Ponce de León & García-Varela, 2017				
<i>M. karlae</i> Pinacho-Pinacho, Hernández-Orts, Sereno-Urbe, Pérez-Ponce de León & García-Varela, 2017	<i>Thorichthys ellioti</i> (Steindachner)	Veracruz, Mexico	KY077066	Pinacho-Pinacho <i>et al.</i> (2017)
Genus: <i>Neoechinorhynchus</i> Stiles & Hassall, 1905				
Subgenus: <i>Neoechinorhynchus</i> Hamann, 1892				
<i>N. (N.) brentnickoli</i> Monks, Pulido-Flores & Violante-Gonzalez, 2011	<i>Dormitator latifrons</i> (Richardson)	Colima, Mexico	KR086199	Pinacho-Pinacho <i>et al.</i> (2015)
<i>N. (N.) bullocki</i> Doolin & Reyda, 2018	<i>Catostomus commersonii</i> (Lacépède)	New York, USA	MK017763	Doolin & Reyda (2018)
<i>N. (N.) chimalapasensis</i> Salgado-Maldonado, 2010	<i>Awaous banana</i> (Valenciennes)	Oaxaca, Mexico	KR086336	Pinacho-Pinacho <i>et al.</i> (2015)
<i>N. (N.) costarricense</i> Pinacho-Pinacho, Sereno-Urbe, Pérez-Ponce de León & García-Varela, 2018	<i>Parachromis managuensis</i> (Günther)	Lago Jalapa, Costa Rica	KR086239 <sup>a</sup>	Pinacho-Pinacho <i>et al.</i> (2015)
<i>N. (N.) cristatus</i> Lynch, 1936	<i>Catostomus macrocheilus</i> Girard	Oregon, USA	MK017786	Doolin & Reyda (2018)
<i>N. (N.) cylindricus</i> (Van Cleave, 1913)	<i>Micropterus salmoides</i> (Lacépède)	Tamaulipas, Mexico	KY077073	Pinacho-Pinacho <i>et al.</i> (2017)
<i>N. (N.) emyditoides</i> Fisher, 1960	<i>Trachemys scripta</i> (Thunberg)	Veracruz, Mexico	HQ634781	García-Varela <i>et al.</i> (2011)
<i>N. (N.) golvani</i> Salgado-Maldonado, 1978	<i>Vieja fenestrata</i> (Günther)	Veracruz, Mexico	FJ388986	Martínez-Aquino <i>et al.</i> (2009)
<i>N. (N.) mamesi</i> Pinacho-Pinacho, Pérez-Ponce de León & García-Varela, 2012	<i>Dormitator latifrons</i> (Richardson)	Chiapas, Mexico	JN830770	Pinacho-Pinacho <i>et al.</i> (2012)
<i>N. (N.) mexicana</i> Pinacho-Pinacho, Sereno-Urbe & García-Varela, 2014	<i>Dormitator maculatus</i> (Bloch)	Veracruz, Mexico	KR086299	Pinacho-Pinacho <i>et al.</i> (2015)

(Continued)

Table 1. (Continued.)

Species	Host	Locality	GenBank accession no.	Reference
<i>N. (N.) panucensis</i> Salgado Maldonado, 2013	<i>Herichthys cyanoguttatus</i> Baird & Girard, 1854	Tamaulipas, Mexico	KY077070	Pinacho-Pinacho et al. (2017)
<i>N. (N.) roseum</i> Salgado-Maldonado, 1978	<i>Citharichthys gilberti</i> Jenkins & Evermann, 1889	Nayarit, Mexico	FJ389000	Martinez-Aquino et al. (2009)
<i>N. (N.) saginatus</i> Van Cleave & Bangham, 1949	na	na	AY829091	García-Varela & Nadler (2005)
<i>N. (N.) schmidti</i> Barger, Thatcher & Nickol, 2004	<i>Trachemys venusta</i> (Gray)	Tabasco, Mexico	FJ389001	Martinez-Aquino et al. (2009)
<b>Outgroups</b>				
<b>Class: Archiacanthocephala</b>				
Order: Moniliformida				
Family: Moniliformidae				
Genus: <i>Moniliformis</i> Travassos, 1915				
<i>M. moniliformis</i> (Bremser, 1811)	<i>Rattus rattus</i> L.	na	AY829086	García-Varela & Nadler (2005)
Order: Oligacanthorhynchida				
Family: Oligacanthorhynchidae				
Genus: <i>Macracanthorhynchus</i> Travassos, 1917				
<i>M. hirudinaceus</i> (Pallas, 1781)	<i>Sus scrofa leucomystax</i> Temminck	Yamaguchi, Japan	LC350000	Kamimura et al. (2018)
Genus: <i>Oligacanthorhynchus</i> Travassos, 1915				
<i>O. tortuosa</i> (Leidy, 1850)	na	na	AY210466	Passamaneck & Halanych (2006)
Class: Palaeacanthocephala				
Order: Echinorhynchida				
Family: Echinorhynchidae				
Genus: <i>Acanthocephalus</i> Koelreuther, 1771				
<i>A. dirus</i> (Van Cleave, 1931)	<i>Asellus aquaticus</i> (L.)	na	AY829106	García-Varela & Nadler (2005)
Family: Heteracanthocephalidae				
Genus: <i>Koronacantha</i> Monks & Pérez-Ponce de León, 1996				
<i>K. mexicana</i> Monks & Pérez-Ponce de León, 1996	<i>Pomadasys leuciscus</i> (Günther)	na	AY829095	García-Varela & Nadler (2005)
Family: Pomphorhynchidae				
Genus: <i>Pomphorhynchus</i> Monticelli, 1905				
<i>P. bullocki</i> Gupta & Lata, 1968	<i>Nasua narica</i> (L.)	na	AY829096	García-Varela & Nadler (2005)
Family: Rhadinorhynchidae				
Genus: <i>Leptorhynchoides</i> Kostylew, 1924				
<i>L. thecatus</i> (Linton, 1891)	<i>Lepomis cyanellus</i> Rafinesque	na	AY829093	García-Varela & Nadler (2005)
Order: Polymorphida				
Family: Polymorphidae				
Genus: <i>Corynosoma</i> Lühe, 1904				
<i>C. hanna</i> Zdzitowiecki, 1984	<i>Phocarctos hookeri</i> (Gray)	New Zealand	JX442180 <sup>b</sup>	García-Varela et al. (2013)
Genus: <i>Hexaglandula</i> Petrochenko, 1950				

(Continued)

Table 1. (Continued.)

Species	Host	Locality	GenBank accession no.	Reference
<i>H. corynosoma</i> (Travassos, 1915)	<i>Nyctanassa violacea</i> (L.)	Veracruz, Mexico	EU267817	García-Varela <i>et al.</i> (2009)
Genus: <i>Pseudocorynosoma</i> Aznar, Pérez-Ponce de León & Raga, 2006				
<i>P. anatarium</i> (Van Cleave, 1945)	<i>Bucephala albeola</i> (L.)	Durango, Mexico	EU267813	García-Varela <i>et al.</i> (2009)
<b>Class: Polyacanthocephala</b>				
Order: Polyacanthorhynchida				
Family: Polyacanthorhynchidae				
Genus: <i>Polyacanthorhynchus</i> Travassos, 1920				
<i>P. caballeroi</i> Diaz-Ungria & Rodrigo, 1960	<i>Caiman yacare</i> Daudin	na	DQ089738	García-Varela & Nadler (2006)

<sup>a</sup>Identified as *Neoechinorhynchus* (*N.*) *golvani* by Pinacho-Pinacho *et al.* (2015) (see Pinacho-Pinacho *et al.*, 2019).

<sup>b</sup>Identified as *Corynosoma australe* Johnston, 1937 by García-Varela *et al.* (2013) (see Hernández-Orts *et al.*, 2017).

subequal, elongate, longer than proboscis receptacle. Gonopore terminal in both sexes (figs 1a, c and 4d, e). Genital spines absent.

**Male.** Based on 18 mounted specimens and four for SEM. Trunk 3.4–7.2 mm (5.3 mm;  $n = 18$ )  $\times$  0.44–1.04 mm (0.7 mm;  $n = 18$ ). Anterior trunk with 14–19 circles of rosethorn-shaped cuticular spines, covering about 14–15% of trunk, reaching to posterior level proboscis receptacle (figs 1b and 3a). Proboscis 130–158 (141;  $n = 5$ )  $\times$  81–107 (96;  $n = 5$ ). Anterior, middle and posterior hooks 49–63 (58;  $n = 15$ ), 43–61 (55;  $n = 12$ ), 34–42 (38;  $n = 9$ ) long, respectively. Roots of anterior, middle and posterior hooks 28–38 (33;  $n = 14$ ), 28–38 (32;  $n = 11$ ), 22–35 (27;  $n = 9$ ) long, respectively. Neck 42–93 (55;  $n = 4$ )  $\times$  127–176 (144;  $n = 4$ ). Proboscis receptacle 341–984 (646;  $n = 10$ )  $\times$  59–121 (93;  $n = 11$ ). Lemnisci 0.90–1.47 mm (1.13 mm;  $n = 8$ ) long. Reproductive system occupying approximately 43–63% (54%;  $n = 11$ ) of trunk length (fig. 1a). Testes equatorial, ovoid, in tandem, often overlapped (fig. 1a). Anterior testis 549–1347 (850;  $n = 11$ )  $\times$  264–625 (430;  $n = 11$ ). Posterior testis 597–1209 (851;  $n = 10$ )  $\times$  266–658 (439;  $n = 10$ ). Cement gland subspherical, 278–612 (454;  $n = 10$ )  $\times$  234–443 (339;  $n = 9$ ), with 4–6 large giant nuclei (fig. 1a). Sperm ducts join seminal vesicle at approximately posterior level of cement gland (fig. 1a). Saeftigen's pouch dorsal to cement duct and seminal vesicle (fig. 1a). Cement duct joins with seminal vesicle duct and Saeftigen's pouch anterior to copulatory bursa. Glandular cellular cluster observed at posterior end of body (fig. 2c). Sperm and mature eggs sometimes observed inside inverted copulatory bursa (fig. 2c). Posterior male reproductive structures extending into fully everted bursa (fig. 2d). Everted copulatory bursa 360–720 (484;  $n = 5$ )  $\times$  210–336 (275;  $n = 5$ ) (figs 2d and 3e). Cement plug observed in some specimens (fig. 3f).

**Female.** Based on 19 gravid mounted specimens and four for SEM. Trunk 7.7–12.9 mm (10.5 mm;  $n = 14$ )  $\times$  0.4–1.2 mm (0.9 mm;  $n = 14$ ). Anterior trunk with 16–19 circles rosethorn-shaped spines (figs 2b and 4c), covering about 5–11% of trunk, reaching almost to posterior level of lemnisci (fig. 1d). Proboscis 114–198 (151;  $n = 7$ )  $\times$  92–134 (112;  $n = 6$ ). Anterior, middle and posterior hooks 57–62 (59;  $n = 11$ ), 53–60 (57;  $n = 7$ ), 33–42 (39;  $n = 9$ ) long, respectively. Roots of anterior, middle and posterior hooks 29–34 (32;  $n = 10$ ), 28–41 (31;  $n = 6$ ), 26–34 (30;  $n = 9$ )

long, respectively. Neck 56–79 (67;  $n = 4$ )  $\times$  138–167 (150;  $n = 4$ ). Proboscis receptacle 572–1031 (801;  $n = 10$ )  $\times$  102–149 (125;  $n = 9$ ). Lemnisci 1.50–1.64 mm (1.56 mm;  $n = 3$ ) long. Reproductive system 1.68–1.79 mm (1.73 mm;  $n = 2$ ) long, post-equatorial, representing 15–16% of trunk length. Uterine bell attached to ventral body wall, with subspherical uterine bell cells (fig. 2e). Uterus differentiated to tubular and muscular part separated by sphincter (fig. 2e). Vaginal bulb, simple, globular, 35–106 (67;  $n = 13$ ) long. Vagina bulb flanked with paired conical muscular jacket (vaginal sleeve), 204–501 (341;  $n = 5$ ) long, extending anteriorly as reproductive ligaments and attaching posteriorly at posterior end of trunk (fig. 2e, f). Posterior end of female sometimes contracted after mating (fig. 4d, e). Mature eggs containing a fully developed acanthor fusiform (fig. 2g), elongate, 24–29 (26;  $n = 32$ )  $\times$  9–11 (10;  $n = 28$ ).

### Taxonomic summary

*Type host.* *Luciobarbus setivimensis* (Valenciennes) (Cypriniformes: Cyprinidae).

*Additional host.* *Luciobarbus callensis* (Valenciennes) (Cypriniformes: Cyprinidae), Algerian barb.

*Type locality.* Azrou (Middle Atlas), Morocco.


*New localities.* Oued Charef dam lake, Souk Ahras City and Zit Emba dam lake, Skikda City, Algeria (no acanthocephalans were collected from Beni Haroun dam lake).

*Site of infection:* Intestine.

*Infection parameters.* Oued Charef dam lake: prevalence 32% ( $n = 289$ ); abundance = 0.9; mean intensity = 2.7; intensity range = 1–12. Zit Emba dam lake: prevalence 36% ( $n = 36$ ); abundance = 1.4; mean intensity = 3.9; intensity range = 1–12.

*Voucher material.* Four males and three females (IPCAS A-109); three males and two females (NHMUK 2019.8.12.1-6); one male and three females (HCMLP-He 7541).

*Representative sequences.* MK953673 (28S rRNA, partial sequence).



**Fig. 1.** *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) ex *Luciobarbus callensis* (Valenciennes) from Oued Charef dam lake, Algeria. (a) Male, whole mount, lateral view. Arrow indicates the posterior end of female inserted through the genital pore of the male. (b) Anterior end of male, lateral view. (c) Female, whole mount, lateral view. (d) Anterior end of female, lateral view. Abbreviations: cg, cephalic ganglion; eg, egg; ghn, giant hypodermal nucleus; le, lemniscus; pr, proboscis receptacle; prs, para-receptacle structure.

### Remarks


The newly collected specimens from *L. callensis* in Algeria are morphologically similar to those described by Dollfus (1951) and Meddour et al. (2010) in having a similar body length, proboscis size, length of the anterior and middle hooks (which are larger than the posterior hooks), shape of roots of the hooks (simple without manubria) and the number of circles of tegumental spines (supplementary table S1). The eggs are also similar in size, although the eggs illustrated in Fig. 6 by Meddour et al. (2010) are smaller ( $28\text{--}30\ \mu\text{m} \times 10\text{--}12\ \mu\text{m}$ ,  $n = 6$ ) than mentioned in their text ( $34\text{--}38\ \mu\text{m} \times 9\text{--}15\ \mu\text{m}$ ).

The newly collected material enlarges the range of intraspecific variability for the number of dorsal hypodermal nuclei, the length of the trunk and the anterior hooks of males, the size of the testis and the size of the eggs, all values being lower than those previously reported in the literature (see supplementary table S1). In contrast, our study also revealed the higher upper limits for the number of circles of cuticular spines, the size of the proboscis of females, the length of the middle hooks for males and females, the length of the posterior hooks of females, the length of the proboscis receptacle of males and the length of lemniscus for males and females (supplementary table S1).


Our morphological observations revealed that the proboscis hooks of *A. (A.) maroccanus* are not arranged in perfect (regular), or nearly perfect, circles reported for this species in the key to the species of the subgenus *Acanthosentis* provided by Amin (2005). Our study has shown that two lateral hooks in the anterior and middle circle are displaced posteriorly. This arrangement of the hooks of *A. (A.) maroccanus* was already observed by Dollfus (1951) and Meddour et al. (2010).

The present study also shows that the posterior male reproductive structures extend into the everted bursa in *A. (A.) maroccanus* (fig. 2d). This feature was previously described only in *A. (A.) parareceptaculis* Amin, 2005 from Japan by Amin (2005), although this characteristic may also be present in *A. (A.) phillipi* (Mashego, 1988) from South Africa (see Fig. 4 in Mashego, 1988). However, as noted by Amin (2005), it is uncertain if this feature appears in other species of the subgenus *Acanthosentis*, mainly because the everted bursa has not been described for some species.

Examination of the newly collected specimens of *A. (A.) maroccanus* has also shown the presence of two structures, which were not reported previously, and which are uncommon among species of the subgenus *Acanthosentis*, i.e. the para-receptacle



**Fig. 2.** *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) ex *Luciobarbus callensis* (Valenciennes) from Oued Charef dam lake, Algeria. (a) Proboscis armature of female, lateral view. (b) Trunk spines of female, lateral view. (c) Posterior end of male with an inverted bursa, showing sperm, mature eggs and glandular cellular cluster at the posterior end of the body. (d) Posterior end of male with detail of reproductive structures extending into the copulatory bursa. (e) Female reproductive system, lateral view. (f) Posterior end of female showing a paired vaginal sleeve. (g) Mature egg. Abbreviations: cd, cement duct; eg, egg; gcc, glandular cellular cluster; ecb, everted copulatory bursa; icb, inverted copulatory bursa; mc, muscular cap of copulatory bursa; mu, muscular uterus; sp, Saeftigen's pouch; spe, sperm; sv, seminal vesicle; tu, tubular uterus; ub, uterine bell; ubc, uterine bell cells; us, uterine sphincter; vb, vagina bulb; vs, vaginal sleeve.




**Fig. 3.** Scanning electron micrographs of adult males of *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) ex *Luciobarbus callensis* (Valenciennes) from Oued Charef dam lake, Algeria. (a) Anterior end, lateral view. (b) Proboscis armature, lateral view. (c) Proboscis, apical view. Arrow indicates the lateral anterior hook displaced posteriorly. (d) Posterior hooks near the base of proboscis, ventral view. (e) Copulatory bursa, lateral view. (f) Copulatory cap, ventral view.

structure and the vaginal sleeve. The para-receptacle structure has also been reported in species of *Neoechinorhynchus*, *Paratenuisentis* Bullock & Samuel, 1975 and *Tenuisentis* Van Cleave, 1936 (see Amin *et al.*, 2007, 2016, 2018; Herlyn & Taraschewski, 2017). To our knowledge, the para-receptacle structure has been observed in both sexes only in *A. (A.) parareceptaclis*, *A. (A.) barmeshoori* Amin, Gholami, Akhlaghi & Heckmann, 2013 and *A. (A.) kashmirensis* Amin, Heckmann & Zargar, 2017. However, *A. (A.) parareceptaclis* and *A. (A.) barmeshoori* are characterized by having two para-receptacle structures, one contiguous with the ventral side of the proboscis receptacle and the other at the posterior end of the trunk (see Amin, 2005; Amin *et al.*, 2013), whereas only the para-receptacle structure contiguous with the ventral side of receptacle is evident in *A. (A.) kashmirensis* (Amin *et al.*, 2017). In *A. (A.) maroccanus*, only the antero-ventral para-receptacle structure was observed (fig. 1b, d), whereas the similar posterior structure is apparently absent (fig. 2c, d), especially in females (fig. 2e, f). A vaginal sleeve was described only in *A. (A.)*

*parareceptaclis* by Amin (2005), but this structure could also be present in *A. (A.) tripathi* Rai, 1967 from India, as indicated by Rai (1967).

During the parasitological examination of Algerian barbs in our study, numerous copulating pairs of *A. (A.) maroccanus* were collected. Both individuals were found firmly attached to the intestinal wall of the fish. The fully everted copulatory bursa of male attaches to the posterior end of the female. Then, the bursa is withdrawn into the body without releasing the female, and the attached portion of the female is inserted through the genital pore of the male (fig. 1a). Inside the invaginated bursa, the muscular cap of the bursa is contracted anteriorly (fig. 1a; see also Fig. 61 in Dollfus, 1951), and the female is firmly trapped by the genital pore of the male (figs 1a and 4f), suggesting that insemination may occur before the female is lodged inside the male. In some copulating females, ovarian balls and unripe eggs were observed, indicating that copulation occurs more than once. Moreover, in one copulating pair, pieces of the copulatory





**Fig. 4.** Scanning electron micrographs of adult females of *Acanthogyrus (Acanthosentis) maroccanus* (Dollfus, 1951) ex *Luciobarbus callensis* (Valenciennes) from Oued Charef dam lake, Algeria. (a) Anterior end, lateral view. (b) Proboscis armature, lateral view. (c) Trunk spines, ventral view. (d) Posterior end, sub-apical view. (e) Genital pore, ventral view. (f) Male and female in copulation. Note that the posterior end of the female (right) is inserted into the posterior end of the male (left).


cap of females were observed in the invaginated bursa of the male. When the copulating pair is separated, the posterior end of the female is strongly contracted (fig. 4d, e), but the posterior end returns to its normal shape after some time (figs 1c and 2e, f). In some separated males, sperm and unripe eggs were observed in the invaginated bursa (fig. 2c). A similar copulatory behaviour was observed in *A. (A.) holospinus* (Sen, 1938), *A. (A.) dattai* (Podder, 1938) and *A. (A.) tilapiae* (see Podder, 1938; Sen, 1938; Baylis, 1947).

### Molecular analyses

Three identical sequences for the D1–D3 region of the 28S rDNA (991–1085 nt long) were generated from isolates of *A. (A.) maroccanus* collected from *L. callensis* from Algeria. The alignment for the 28S rDNA dataset comprised 1618 nt positions, of which 599 (37%) were parsimony-informative. Phylogenetic relations

estimated by maximum likelihood and Bayesian inference methods resulted in consensus trees with identical topologies (Fig. 5).

The phylogenetic relationships at the class level were very similar to the phylogenetic hypotheses obtained in previous studies (e.g. García-Varela & Nadler, 2005; Verweyen *et al.*, 2011), where the Polyacanthocephala was the sister clade of the Eoacanthocephala. Within the Eoacanthocephala, which includes members of the Gyraacanthocephala and the Neoechinorhynchida, our sequence of *A. (A.) maroccanus* was placed in a well-supported clade with *P. (D.) ophiocephalus*, both species included in the Quadrigyridae, the only family within the Gyraacanthocephala. The clade formed by the members of the Gyraacanthocephala was found to be sister of a clade including all members of the Neoechinorhynchida, but with low support. Within the Neoechinorhynchida, the phylogenetic analyses showed *Neoechinorhynchus (N.) roseum* (GenBank accession no. FJ389000) as an earliest diverging lineage and two further groups. The first well-supported group was formed by two clades



**Fig. 5.** Phylogenetic tree resulting from maximum likelihood analysis of partial (D1–D3 regions) sequences of 28S rDNA of *Acanthogyrus (Acanthosentis) maroccanus* from Algeria. Nodal supports are depicted as bootstraps values ( $\geq 70\%$  shown only) followed by posterior probabilities ( $\geq 0.70$  shown only). The branch-length scale bar indicates the expected number of substitutions per site. The newly generated sequence is indicated in bold.

of *Neoechinorhynchus* spp. (the isolates of three and four species, respectively) and *Mayarhynchus karlae* (GenBank accession no. KY077066). The second group, having low support, was formed by a clade of *Floridosentis mugilis* (GenBank accession no. JQ436495), *F. pacifica* (GenBank accession no. JQ436531) and *A. duranguensis* (GenBank accession no. KY077080), and its sister clade of six species of *Neoechinorhynchus*. Under the current taxon sampling, *Neoechinorhynchus* was resolved as a polyphyletic group with at least three separate clades and the lineage of *N. (N.) roseum*. The present phylogenetic analysis is largely consistent with previous phylogenetic assessments for species of *Neoechinorhynchus* (e.g. García-Varela & Pinacho-Pinacho, 2019).

The genetic divergence value between *A. (A.) maroccanus* and *P. (D.) ophiocephalus* (the other available member of the Quadrigyridae) was 39.2%. Variation of the interspecific genetic divergence between *A. (A.) maroccanus* and other members of the Eoacanthocephala ranged between 33.9% (*Neoechinorhynchus (Neoechinorhynchus) golvani*) and 43.6% (*F. pacifica*).

## Discussion

*Acanthogyrus (A.) maroccanus* has been registered only in *L. setivimensis* and *L. callensis* from Algeria and Morocco (Dollfus, 1951; Meddour, 2009; Meddour et al., 2010). This acanthocephalan has been reported in freshwater cyprinids collected in Oubeira Lake and several dam lakes (i.e. Ain Dalia, Bouhamdane, Oued Charef and Zit Emba) or streams (i.e. Bounamoussa, El Kebir and Seybouse) from East Algeria (Meddour et al., 2010; present study).

The interspecific relationships and phylogenetic position of *Acanthogyrus* remain poorly explored. According to the GenBank dataset, molecular data for mitochondrial and nuclear genes have been provided only for two of the 47 species of the genus. Chenuil et al. (1997) provided a partial fragment of the 28S rRNA gene of *A. (A.) tilapiae*; however, this is extremely short (311 bp) and, thus, could hardly be compared with our newly generated sequences, which are much longer (991–

1085 bp). Even though our phylogenetic analysis showed that our isolate of *A. (A.) maroccanus* forms a poorly supported clade within the Eoacanthocephala, the topology recovered from different analyses was always identical. The phylogenetic signal detected in our study shows the limitation of the currently available molecular data to understand the interrelations of species of the Quadrigyridae.

**Supplementary material.** To view supplementary material for this article, please visit <https://doi.org/10.1017/S0022149X19000737>.

**Acknowledgements.** The authors are indebted to two anonymous reviewers for their helpful suggestions, and to M. Borovková and R. Kuchta, IPCAS, České Budějovice for their technical help during the examination of the acanthocephalans. Thanks are also due to D. Hernández-Mena, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional – Unidad Mérida, Yucatán, Mexico for his valuable comments and assistance with the phylogenetic analyses.

**Financial support.** This study was partly supported by the Faculty of Natural and Life Sciences, University of Mohamed Cherif Messaadia of Algeria (CNEPRU: 410120150001), Institute of Parasitology, Biology Centre of the Czech Academy of Sciences (RVO: 60077344), Czech Science Foundation (project No. P505/12/G112), Grant Agency VEGA (project No. 2/0159/16), and Agencia Nacional de Promoción Científica y Tecnológica of Argentina (PICT No. 2017-3232).

**Conflicts of interest.** None.

**Ethical standards.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional guides on the care and use of animals.

## References

- Amin OM (2005) Occurrence of the subgenus *Acanthosentis* Verma & Datta, 1929 (Acanthocephala: Quadrigyridae) in Japan, with the description of *Acanthogyrus (Acanthosentis) alternatospinus* n. sp. and *A. (A.) parareceptaculis* n. sp. from Lake Biwa drainage fishes and a key to the species of the subgenus. *Systematic Parasitology* **60**, 125–137.

- Amin OM** (2013) Classification of the Acanthocephala. *Folia Parasitologica* **60**, 273–305.
- Amin OM and Hendrix SS** (1999) Acanthocephala of cichlids (Pisces) in Lake Malawi, Africa, with a description of *Acanthogyrus (Acanthosentis) malawiensis* sp. n. (Quadrigyridae) from *Labeo cylindricus* Peters, 1852 (Cyprinidae). *Journal of the Helminthological Society of Washington* **66**, 47–55.
- Amin OM, Heckmann R and Standing MD** (2007) The structural-functional relationship of the para-receptacle structure in Acanthocephala. *Comparative Parasitology* **74**, 383–387.
- Amin OM, Heckmann RA and Zargar UR** (2017) Description of a new quadrigyrid acanthocephalan from Kashmir, with notes on metal analysis and histopathology, and a key to species of the subgenus *Acanthosentis* from the Indian subcontinent. *Journal of Parasitology* **103**, 458–470.
- Amin OM, Heckmann RA and Ha NV** (2018) Descriptions of *Acanthocephalus parallelelementglandatus* (Echinorhynchidae) and *Neoechinorhynchus (N.) pennahia* (Neoechinorhynchidae) (Acanthocephala) from amphibians and fish in Central and Pacific coast of Vietnam, with notes on *N. (N.) longnucleatus*. *Acta Parasitologica* **63**, 572–585.
- Amin OM, Gholami Z, Akhlaghi M, et al.** (2013) The description and host-parasite relationships of a new quadrigyrid species (Acanthocephala) from the Persian tooth-carp, *Aphanius farsicus* (Actinopterygii: Cyprinodontidae) in Iran. *Journal of Parasitology* **99**, 257–263.
- Amin OM, Evans RP, Boungou M, et al.** (2016) Morphological and molecular description of *Tenuisentis niloticus* (Meyer, 1932) (Acanthocephala: Tenuisentidae) from *Heterotis niloticus* (Cuvier) (Actinopterygii: Arapaimidae), in Burkina Faso, with emendation of the family diagnosis and notes on new features, cryptic genetic diversity and histopathology. *Systematic Parasitology* **93**, 173–191.
- Baylis HA** (1947) A new acanthocephalan from an East African freshwater fish. *Annals and Magazine of Natural History* **14**, 861–868.
- Bush AO, Lafferty K, Lotz J, et al.** (1997) Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology* **83**, 575–583.
- Chenuil A, Solignac M and Bernard M** (1997) Evolution of the large-subunit ribosomal RNA binding site for protein L23/25. *Molecular Biology and Evolution* **14**, 578–588.
- Darriba D, Taboada GL, Doallo R, et al.** (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* **9**, 772.
- Dollfus RPH** (1951) Miscellanea helminthologica Marocana. III. Quelques trématodes, cestodes et acanthocéphales. *Archives de l'Institut Pasteur du Maroc* **4**, 104–229.
- Doolin ML and Reyda FB** (2018) A new species of *Neoechinorhynchus* (Acanthocephala: Neoechinorhynchidae) from white sucker (*Catostomus commersonii*) in New York. *Journal of Parasitology* **104**, 671–678.
- García-Varela M and Nadler SA** (2005) Phylogenetic relationships of Palaeacanthocephala (Acanthocephala) inferred from SSU and LSU rDNA gene sequences. *Journal of Parasitology* **91**, 1401–1409.
- García-Varela M and Nadler SA** (2006) Phylogenetic relationships among Syndermata inferred from nuclear and mitochondrial gene sequences. *Molecular Phylogenetics and Evolution* **40**, 61–72.
- García-Varela M and Pinacho-Pinacho CD** (2019) Molecular characterization of *Neoechinorhynchus cylindratus* Van Cleave, 1913 (Acanthocephala: Neoechinorhynchidae), a parasite of the largemouth bass (*Micropterus salmoides*) in northern Mexico. *Journal of Helminthology*. <https://doi.org/10.1017/S0022149X18001104>
- García-Varela M, García-Prieto L and Rodríguez RP** (2011) Molecular identification and first description of the male of *Neoechinorhynchus schmidti* (Acanthocephala: Neoechinorhynchidae), a parasite of *Trachemys scripta* (Testudines) in Mexico. *Parasitology International* **60**, 433–439.
- García-Varela M, Pérez-Ponce de León G, Aznar FJ, et al.** (2009) Systematic position of *Pseudocorynosoma* and *Andracantha* (Acanthocephala, Polymorphidae) based on nuclear and mitochondrial gene sequences. *Journal of Parasitology* **95**, 178–185.
- García-Varela M, Pérez-Ponce de León G, Aznar FJ and Nadler SA** (2013) Phylogenetic relationship among genera of Polymorphidae (Acanthocephala), inferred from nuclear and mitochondrial gene sequences. *Molecular Phylogenetics and Evolution* **68**, 176–184.
- Herlyn H and Taraschewski H** (2017) Evolutionary anatomy of the muscular apparatus involved in the anchoring of Acanthocephala to the intestinal wall of their vertebrate hosts. *Parasitology Research* **116**, 1207–1225.
- Hernández-Orts JS, Smales LR, Pinacho-Pinacho CD, et al.** (2017) Novel morphological and molecular data for *Corynosoma hanna* Zdzitowiecki, 1984 (Acanthocephala: Polymorphidae) from teleosts, fish-eating birds and pinnipeds from New Zealand. *Parasitology International* **66**, 905–916.
- Huelsenbeck JP and Ronquist F** (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* **17**, 754–755.
- Kamimura K, Yonemitsu K, Maeda K, et al.** (2018) An unexpected case of a Japanese wild boar (*Sus scrofa leucomystax*) infected with the giant thorny-headed worm (*Macracanthorhynchus hirudinaceus*) on the mainland of Japan (Honshu). *Parasitology Research* **117**, 2315–2322.
- Kearse M, Moir R, Wilson A, et al.** (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649.
- Littlewood DTJ, Curini-Galletti M and Herniou EA** (2000) The interrelationships of Proseriata (Platyhelminthes: Seriata) flatworms tested with molecules and morphology. *Molecular Phylogenetics and Evolution* **16**, 449–466.
- Martínez-Aquino A, Reyna-Fabián ME, Rosas-Valdez R, et al.** (2009) Detecting a complex of cryptic species within *Neoechinorhynchus golvani* (Acanthocephala: Neoechinorhynchidae) inferred from ITSs and LSU rDNA gene sequences. *Journal of Parasitology* **95**, 1040–1047.
- Mashego SN** (1988) A new species of *Acanthosentis* Verma & Datta, 1929 (Acanthocephala: Quadrigyridae) from *Barbus neefi* in South Africa. *Annals of the Transvaal Museum* **34**, 545–549.
- Meddour A** (2009) *Pisciculture et biodiversité de la parasitofaune des poissons dans le Nord-Est de l'Algérie*. 236 pp. PhD Thesis, El Taref, University of El Taref.
- Meddour A, Meddour-Bouderda K, Brahim-Tazi NA, et al.** (2010) Microscopie électronique à balayage des parasites des poissons du lac Oubeira Algérie. *European Journal of Scientific Research* **48**, 129–141.
- Olson PD, Cribb TH, Tkach VV, et al.** (2003) Phylogeny and classification of the Digenea (Platyhelminthes: Trematoda). *International Journal for Parasitology* **33**, 733–755.
- Passamanek Y and Halanych KM** (2006) Lophotrochozoan phylogeny assessed with LSU and SSU data: Evidence of lophophorate polyphyly. *Molecular Phylogenetics and Evolution* **40**, 20–28.
- Pinacho-Pinacho CD, Pérez-Ponce de León G and García-Varela M** (2012) Description of a new species of *Neoechinorhynchus* (Acanthocephala: Neoechinorhynchidae) a parasite of *Dormitator latifrons* from Southwestern Mexico based on morphological and molecular characters. *Parasitology International* **61**, 634–644.
- Pinacho-Pinacho CD, Sereno-Urbe AL, Pérez-Ponce de León G, et al.** (2015) Checklist of the species of *Neoechinorhynchus* (Acanthocephala: Neoechinorhynchidae) in fishes and turtles in Middle-America, and their delimitation based on sequences of the 28S rDNA. *Zootaxa* **3985**, 98–116.
- Pinacho-Pinacho CD, Hernández-Orts JS, Sereno-Urbe AL, et al.** (2017) *Mayarhynchus karlae* n. g., n. sp. (Acanthocephala: Neoechinorhynchidae), a parasite of cichlids (Perciformes: Cichlidae) in southeastern Mexico, with comments on the paraphyly of *Neoechinorhynchus* Stiles & Hassall, 1905. *Systematic Parasitology* **94**, 351–365.
- Pinacho-Pinacho CD, Sereno-Urbe AL, García-Varela M, et al.** (2019) A closer look at the morphological and molecular diversity of *Neoechinorhynchus* (Acanthocephala) in Middle American cichlids (Osteichthyes: Cichlidae), with the description of a new species from Costa Rica. *Journal of Helminthology*. <https://doi.org/10.1017/S0022149X18001141>
- Podder TN** (1938) A new species of Acanthocephala, *Acanthosentis dattai*, n. sp., from a fresh-water fish of Bengal, *Barbus ticto* (Ham. & Buch.) and *B. stigma* (Cuv. & Val.). *Parasitology* **30**, 171–175.
- Rai P** (1967) On four acanthocephalan genera parasitic in freshwater fishes with description of three new species. *Indian Journal of Helminthology* **19**, 27–44.
- Rosas-Valdez R, Morrone JJ and García-Varela M** (2012) Molecular phylogenetics of *Floridosentis* Ward, 1953 (Acanthocephala: Neoechinorhynchidae) parasites of mullets (Osteichthyes) from Mexico, using 28S rDNA sequences. *Journal of Parasitology* **98**, 855–862.

- Sen P** (1938) On a new species of Acanthocephala, *Acanthosentis holospinus* sp. nov. from the fish *Barbus stigma* (Cuv. and Val.). *Proceedings of the Indian Academy of Sciences* **7**, 41–46.
- Stamatakis A** (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313.
- Swofford D.L.** (2002) *PAUP\*: Phylogenetic Analysis Using Parsimony (\*and Other Methods)*, 4.0. Sunderland, Massachusetts, Sinauer Associates.
- Thompson JD, Higgins HG and Gibson TJ** (1994) CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**, 4673–4680.
- Verweyen L, Klimpel S and Palm HW** (2011) Molecular phylogeny of the Acanthocephala (class Palaeacanthocephala) with a paraphyletic assemblage of the orders Polymorphida and Echinorhynchida. *PLoS ONE* **6**, e28285.
- Werle E, Schneider C, Renner M, et al.** (1994) Convenient single-step, one tube purification of PCR products for direct sequencing. *Nucleic Acids Research* **22**, 4354–4355.